

BICOMMUTANTS OF REDUCED UNBOUNDED OPERATOR ALGEBRAS

FABIO BAGARELLO, ATSUSHI INOUE, AND CAMILLO TRAPANI

ABSTRACT. The unbounded bicommutant $(\mathfrak{M}_{E'})''_{\text{wc}}$ of the *reduction* of an O^* -algebra $\mathfrak{M}$ via a given projection E' weakly commuting with $\mathfrak{M}$ is studied, with the aim of finding conditions under which the reduction of a GW^* -algebra is a GW^* -algebra itself. The obtained results are applied to the problem of the existence of conditional expectations on O^* -algebras.

1. INTRODUCTION AND PRELIMINARIES

If $\mathfrak{M}$ is a von Neumann algebra and E' a projection on the commutant $\mathfrak{M}'$ of $\mathfrak{M}$, then it is well known that the *reduced* algebra $\mathfrak{M}_{E'} = \{XE'; X \in \mathfrak{M}\}$ is again a von Neumann algebra. The notion of von Neumann algebra has been generalized to unbounded operator algebras by introducing several classes of O^* -algebras, such as GW^* -algebras, EW^* -algebras, etc. It is then natural and important for applications to pose the question if the *reduced* algebra of a GW^* -algebra via a projection E' picked in its weak bounded commutant is again a GW^* -algebra. In order to answer this question we undertake a more general study of the reduction process for O^* -algebras. We begin with reviewing the basic definitions and properties of O^* -algebras needed in this paper and refer to [5, 6] for more details.

Let $\mathcal{H}$ be a Hilbert space with inner product $\langle \cdot | \cdot \rangle$ and $\mathcal{D}$ a dense subspace of $\mathcal{H}$. We denote by $\mathcal{L}^\dagger(\mathcal{D})$ the set of all linear operators X defined in $\mathcal{D}$ such that $X\mathcal{D} \subset \mathcal{D}$, the domain $D(X^*)$ of its adjoint contains $\mathcal{D}$ and $X^*\mathcal{D} \subset \mathcal{D}$. Then $\mathcal{L}^\dagger(\mathcal{D})$ is a $*$ -algebra with the usual operations: $X+Y$, αX , XY and the involution $X \mapsto X^\dagger := X^* \upharpoonright \mathcal{D}$. A $*$ -subalgebra of $\mathcal{L}^\dagger(\mathcal{D})$ is called an O^* -algebra on $\mathcal{D}$. In this paper we will assume that an O^* -algebra always contains the identity I .

Let $\mathfrak{M}$ be an O^* -algebra on $\mathcal{D}$. The *graph topology* $t_{\mathfrak{M}}$ on $\mathcal{D}$ is the locally convex topology defined by the family $\{\|\cdot\|_X; X \in \mathfrak{M}\}$ of seminorms: $\|\xi\|_X = \|X\xi\|$, $\xi \in \mathcal{D}$. If the locally convex space $\mathcal{D}[t_{\mathfrak{M}}]$ is complete, then $\mathfrak{M}$ is said to be *closed*. In particular, if $\mathcal{D}[t_{\mathfrak{M}}]$ is a Fréchet space, then $\mathcal{D}$ is called the Fréchet domain of $\mathfrak{M}$. More in general, we denote by $\tilde{\mathcal{D}}(\mathfrak{M})$ the completion of the locally convex space $\mathcal{D}[t_{\mathfrak{M}}]$ and put

$$\tilde{X} := \overline{X} \upharpoonright \tilde{\mathcal{D}}(\mathfrak{M}) \quad \text{and} \quad \tilde{\mathfrak{M}} := \{\tilde{X} : X \in \mathfrak{M}\}.$$

Then $\tilde{\mathcal{D}}(\mathfrak{M}) = \bigcap_{X \in \mathfrak{M}} D(\overline{X})$ and $\tilde{\mathfrak{M}}$ is a closed O^* -algebra on $\tilde{\mathcal{D}}(\mathfrak{M})$ which is called the *closure* of $\mathfrak{M}$, since it is the smallest closed extension of $\mathfrak{M}$. We next recall the notion of self-adjointness of $\mathfrak{M}$. If $\mathcal{D} = \mathcal{D}^*(\mathfrak{M}) := \bigcap_{X \in \mathfrak{M}} D(X^*)$, then $\mathfrak{M}$ is said to be *self-adjoint*. If $\tilde{\mathcal{D}}(\mathfrak{M}) = \mathcal{D}^*(\mathfrak{M})$, then $\mathfrak{M}$ is said to be *essentially self-adjoint*. It

2000 *Mathematics Subject Classification.* 47L60.

This work was supported by CORI, Università di Palermo.

is clear that

$$\begin{aligned}\mathcal{D} &\subset \widetilde{\mathcal{D}}(\mathfrak{M}) \subset \mathcal{D}^*(\mathfrak{M}) \\ X &\subset \widetilde{X} \subset X^{\dagger*}, \quad \forall X \in \mathfrak{M}.\end{aligned}$$

The *weak commutant* $\mathfrak{M}'_{\mathfrak{w}}$ of $\mathfrak{M}$ is defined by

$$\mathfrak{M}'_{\mathfrak{w}} = \{C \in \mathcal{B}(\mathcal{H}) : \langle CX\xi | \eta \rangle = \langle C\xi | X^{\dagger}\eta \rangle, \forall X \in \mathfrak{M}, \forall \xi, \eta \in \mathcal{D}\},$$

where $\mathcal{B}(\mathcal{H})$ is the $*$ -algebra of all bounded linear operators on $\mathcal{H}$. Then $\mathfrak{M}'_{\mathfrak{w}}$ is a weak-operator closed $*$ -invariant subspace of $\mathcal{B}(\mathcal{H})$ (but it is not, in general, a von Neumann algebra) and $(\widetilde{\mathfrak{M}})'_{\mathfrak{w}} = \mathfrak{M}'_{\mathfrak{w}}$. If $\mathfrak{M}'_{\mathfrak{w}}\mathcal{D} \subset \mathcal{D}$, as it happens for self-adjoint $\mathfrak{M}$, then $\mathfrak{M}'_{\mathfrak{w}}$ is a von Neumann algebra. In this paper we will use the following bicommutants of $\mathfrak{M}$:

$$(\mathfrak{M}'_{\mathfrak{w}})' = \{A \in \mathcal{B}(\mathcal{H}) : AC = CA, \forall C \in \mathfrak{M}'_{\mathfrak{w}}\},$$

$$\mathfrak{M}''_{\text{wc}} := (\mathfrak{M}'_{\mathfrak{w}})'_{\mathfrak{c}} = \{X \in \mathcal{L}^{\dagger}(\mathcal{D}) : \langle CX\xi | \eta \rangle = \langle C\xi | X^{\dagger}\eta \rangle, \forall C \in \mathfrak{M}'_{\mathfrak{w}}, \forall \xi, \eta \in \mathcal{D}\}.$$

Then, $(\mathfrak{M}'_{\mathfrak{w}})'$ is a von Neumann algebra on $\mathcal{H}$ and $\mathfrak{M}''_{\text{wc}}$ is a τ_{s^*} -closed O^* -algebra on $\mathcal{D}$ such that $\mathfrak{M} \subset \mathfrak{M}''_{\text{wc}}$ and $(\mathfrak{M}''_{\text{wc}})'_{\mathfrak{w}} = \mathfrak{M}'_{\mathfrak{w}}$, where the strong $*$ -topology τ_{s^*} is defined by the family $\{p_{\xi}^*(\cdot); \xi \in \mathcal{D}\}$ of seminorms:

$$p_{\xi}^* := \|X\xi\| + \|X^{\dagger}\xi\|, \quad X \in \mathcal{L}^{\dagger}(\mathcal{D}).$$

Furthermore, if $\mathfrak{M}'_{\mathfrak{w}}\mathcal{D} \subset \mathcal{D}$, then

$$\mathfrak{M}''_{\text{wc}} = \{X \in \mathcal{L}^{\dagger}(\mathcal{D}) : \overline{X} \text{ is affiliated with } (\mathfrak{M}'_{\mathfrak{w}})'\} = \overline{(\mathfrak{M}'_{\mathfrak{w}})' \upharpoonright \mathcal{D}}^{\tau_{s^*}} \cap \mathcal{L}^{\dagger}(\mathcal{D}).$$

An O^* -algebra $\mathfrak{M}$ on $\mathcal{D}$ is called a *GW $*$ -algebra* if $\mathfrak{M}'_{\mathfrak{w}}\mathcal{D} \subset \mathcal{D}$ and $\mathfrak{M} = \mathfrak{M}''_{\text{wc}}$.

Let $\mathfrak{M}$ be a closed O^* -algebra on $\mathcal{D}$ such that $\mathfrak{M}'_{\mathfrak{w}}\mathcal{D} \subset \mathcal{D}$ and let E' be a projection in the von Neumann algebra $\mathfrak{M}'_{\mathfrak{w}}$. Let $\mathfrak{M}_{E'}$ denote the *reduced algebra* $\mathfrak{M}_{E'} = \{XE' : X \in \mathfrak{M}\}$. Then it is known [1] that

$$\begin{aligned}(\mathfrak{M}_{E'})'_{\mathfrak{w}} &= (\mathfrak{M}'_{\mathfrak{w}})_{E'}, \\ ((\mathfrak{M}_{E'})'_{\mathfrak{w}})' &= ((\mathfrak{M}'_{\mathfrak{w}})')_{E'}.\end{aligned}$$

But the following question is open

Question: Does the equality $(\mathfrak{M}_{E'})''_{\text{wc}} = (\mathfrak{M}''_{\text{wc}})_{E'}$ hold?

If the answer to this question is affirmative, then the reduced algebra $\mathfrak{M}_{E'}$ of a given GW $*$ -algebra is again a GW $*$ -algebra. This question will be considered in Section 2, where we will show that if $\mathcal{D}$ is the Fréchet domain of $\mathfrak{M}$ and the linear span $\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle$ of the $\mathfrak{M}$ -invariant subspace $\mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D}$ is essentially self-adjoint, then $(\mathfrak{M}_{E'})''_{\text{wc}} = (\mathfrak{M}''_{\text{wc}})_{E'}$. An $\mathfrak{M}$ -invariant subspace $\mathcal{M}$ of $\mathcal{D}$ is called *essentially self-adjoint* if the O^* -algebra $\mathfrak{M} \upharpoonright \mathcal{M} := \{X \upharpoonright \mathcal{M} : X \in \mathfrak{M}\}$ of $\mathcal{M}$ is essentially selfadjoint.

Furthermore, we will show that, if the graph topology $t_{\mathfrak{M}}$ is defined by a sequence $\{T_n\}$ of essentially self-adjoint operators of $\mathfrak{M}$, whose spectral projections leave the domain $\mathcal{D}$ invariant, then again $(\mathfrak{M}_{E'})''_{\text{wc}} = (\mathfrak{M}''_{\text{wc}})_{E'}$.

In Section 3 we shall apply the results of Section 2 to the study of conditional expectations for O^* -algebras.

2. MAIN RESULTS

Let $\mathfrak{M}$ be a closed O^* -algebra on $\mathcal{D}$ such that $\mathfrak{M}'_{\mathfrak{w}}\mathcal{D} \subset \mathcal{D}$ and let E' be a projection in the von Neumann algebra $\mathfrak{M}'_{\mathfrak{w}}$. In this Section we look for conditions for the equality $(\mathfrak{M}''_{\mathfrak{w}c})_{E'} = (\mathfrak{M}_{E'})''_{\mathfrak{w}c}$ to hold. Since $(\mathfrak{M}_{E'})'_{\mathfrak{w}} = (\mathfrak{M}'_{\mathfrak{w}})_{E'}$, it follows that

$$(1) \quad (\mathfrak{M}''_{\mathfrak{w}c})_{E'} \subset ((\mathfrak{M}'_{\mathfrak{w}})_{E'})'_{\mathfrak{c}} = (\mathfrak{M}_{E'})''_{\mathfrak{w}c}.$$

Hence we need only to prove the converse inclusion: $(\mathfrak{M}_{E'})''_{\mathfrak{w}c} \subset (\mathfrak{M}''_{\mathfrak{w}c})_{E'}$. An element X of $(\mathfrak{M}_{E'})''_{\mathfrak{w}c}$ is an operator on $E'\mathcal{D}$, and so to show that $X \in (\mathfrak{M}''_{\mathfrak{w}c})_{E'}$ we need to extend X to an operator on $\mathcal{D}$. We will consider this extension problem below.

Let Z denote the *central support* of E' i.e., Z is the projection onto the closure of the subspace $\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{H} \rangle$. We define $\mathcal{E} := \langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle \oplus (I - Z)\mathcal{D}$. Then $\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle \subset Z\mathcal{D}$ and $\mathcal{E} \subset \mathcal{D}$.

For $X \in (\mathfrak{M}_{E'})''_{\mathfrak{w}c}$, we put

$$X_e \left(\sum_k C_k E' \xi_k + (I - Z)\eta \right) := \sum_k C_k X E' \xi_k, \quad C_k \in \mathfrak{M}'_{\mathfrak{w}}, \xi_k, \eta \in \mathcal{D}.$$

Then we have the following

Lemma 2.1. $X_e \in \mathcal{L}^\dagger(\mathcal{E})$ and $((\mathfrak{M}_{E'})''_{\mathfrak{w}c})_e := \{X_e : X \in (\mathfrak{M}_{E'})''_{\mathfrak{w}c}\}$ is an O^* -algebra on $\mathcal{E}$.

Proof. For any $X \in (\mathfrak{M}_{E'})''_{\mathfrak{w}c}$, $C_k, K_j \in \mathfrak{M}'_{\mathfrak{w}}$ and $\xi_k, \zeta_j, \eta, \eta_1 \in \mathcal{D}$ we have

$$\begin{aligned} & \left\langle X_e \left(\sum_k C_k E' \xi_k + (I - Z)\eta \right) \middle| \sum_j K_j E' \zeta_j + (I - Z)\eta_1 \right\rangle \\ &= \sum_{k,j} \langle C_k X E' \xi_k \mid K_j E' \zeta_j \rangle \\ &= \sum_{k,j} \langle E' K_j^* C_k E' X E' \xi_k \mid E' \zeta_j \rangle \\ &= \sum_{k,j} \langle E' K_j^* C_k E' \xi_k \mid X^\dagger E' \zeta_j \rangle \\ &= \left\langle \sum_k C_k E' \xi_k \middle| \sum_j K_j X^\dagger E' \zeta_j \right\rangle \\ &= \left\langle \sum_k C_k E' \xi_k + (I - Z)\eta \middle| (X_e)^\dagger \left(\sum_j K_j E' \zeta_j + (I - Z)\eta_1 \right) \right\rangle \end{aligned}$$

which implies that X_e is well-defined and $(X_e)^\dagger = (X^\dagger)_e$. It is clear that X_e is a linear operator on $\mathcal{E}$. Hence $X_e \in \mathcal{L}^\dagger(\mathcal{E})$. Moreover, as it is easily seen, $(XY)_e = X_e Y_e$. Hence, $e(\mathfrak{M}_{E'}) := ((\mathfrak{M}_{E'})''_{\mathfrak{w}c})_e$ is an O^* -algebra on $\mathcal{E}$. $\square$

Since $e(\mathfrak{M}_{E'})$ is an O^* -algebra on $\mathcal{E}$, its closure is an O^* -algebra on $\tilde{\mathcal{E}}$. It is easily seen that $\tilde{\mathcal{E}} = \overline{\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle}^{t_{e(\mathfrak{M}_{E'})}} \oplus (I - Z)\mathcal{D}$. On the other hand, $\mathcal{D} = Z\mathcal{D} \oplus (I - Z)\mathcal{D}$; hence, if $\overline{\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle}^{t_{e(\mathfrak{M}_{E'})}} = Z\mathcal{D}$, we can extend any $X \in (\mathfrak{M}_{E'})''_{\mathfrak{w}c}$ to $\mathcal{D}$.

Theorem 2.2. *Let $\mathfrak{M}$ be a closed O^* -algebra on $\mathcal{D}$ such that $\mathfrak{M}'_{\mathfrak{w}}\mathcal{D} \subset \mathcal{D}$ and let E' be a projection in the von Neumann algebra $\mathfrak{M}'_{\mathfrak{w}}$. Suppose that $\widetilde{\mathcal{E}}((\mathfrak{M}'_{E'})''_{\mathfrak{w}c})_e = \mathcal{D}$ or, equivalently, $\overline{\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle}^{t_{e(\mathfrak{M}_{E'})}} = Z\mathcal{D}$. Then, $(\mathfrak{M}''_{\mathfrak{w}c})_{E'} = (\mathfrak{M}_{E'})''_{\mathfrak{w}c}$.*

Proof. By the assumption, for every $X \in (\mathfrak{M}_{E'})''_{\mathfrak{w}c}$, X_e extends to an operator $\widetilde{X}_e$ on $\mathcal{D}$; that is $\widetilde{X}_e = \overline{X_e}|_{\mathcal{D}}$ and $(\widetilde{X}_e)_{E'} = X$. Furthermore, we have $\widetilde{X}_e \in \mathfrak{M}''_{\mathfrak{w}c}$. Indeed, for every $C \in \mathfrak{M}'_{\mathfrak{w}}$, we have

$$\begin{aligned} CX_e \left(\sum_k C_k E' \xi_k + (I - Z)\eta \right) &= C \sum_k C_k X E' \xi_k \\ &= \sum_k C C_k X E' \xi_k \\ &= X_e C \left(\sum_k C_k E' \xi_k + (I - Z)\eta \right), \end{aligned}$$

for every $C_k \in \mathfrak{M}'_{\mathfrak{w}}$, $\xi_k, \eta \in \mathcal{D}$. Hence, $CX_e = X_e C$ on $\mathcal{E}$ and so $C\widetilde{X}_e = \widetilde{X}_e C$. Thus $\widetilde{X}_e \in \mathfrak{M}''_{\mathfrak{w}c}$ and $X = (\widetilde{X}_e)_{E'} \in (\mathfrak{M}''_{\mathfrak{w}c})_{E'}$. In conclusion, $(\mathfrak{M}_{E'})''_{\mathfrak{w}c} \subset (\mathfrak{M}''_{\mathfrak{w}c})_{E'}$ and by (1) the equality follows. $\square$

The next step consists, of course, in looking for situations where the conditions of Theorem 2.2 are satisfied. We begin with the following

Lemma 2.3. *Let $\mathfrak{M}$ and E' be as above. Suppose that*

(i) *for every $X \in (\mathfrak{M}_{E'})''_{\mathfrak{w}c}$ there exists an element Y of $\mathfrak{M}''_{\mathfrak{w}c}$ such that*

$$\|X_e \xi\| \leq \|Y \xi\|, \quad \forall \xi \in \mathcal{E}$$

(equivalently, $X_e^\dagger X_e \leq Y^\dagger Y$ on $\mathcal{E}$)

(ii) $\overline{\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle}^{t_{\mathfrak{M}''_{\mathfrak{w}c}}} = Z\mathcal{D}$.

Then, $(\mathfrak{M}_{E'})''_{\mathfrak{w}c} = (\mathfrak{M}''_{\mathfrak{w}c})_{E'}$.

Proof. The conditions (i) and (ii) immediately imply that

$$Z\mathcal{D} \subset \overline{\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle}^{t_{e(\mathfrak{M}_{E'})}}.$$

Hence

$$\overline{\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle}^{t_{e(\mathfrak{M}_{E'})}} = Z\mathcal{D}.$$

By Theorem 2.2, we get $(\mathfrak{M}_{E'})''_{\mathfrak{w}c} = (\mathfrak{M}''_{\mathfrak{w}c})_{E'}$. $\square$

Remark 2.4. A comment is in order for condition (ii) of Lemma 2.3. Let us in fact consider the following statements:

(i) The $\mathfrak{M}''_{\mathfrak{w}c}$ -invariant subspace $\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle$ is essentially self-adjoint for $\mathfrak{M}''_{\mathfrak{w}c}$.

(ii) $\overline{\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle}^{t_{\mathfrak{M}''_{\mathfrak{w}c}}} = Z\mathcal{D}$.

Then (i) $\Rightarrow$ (ii). In particular if $\mathfrak{M}$ is self-adjoint, then the two conditions are equivalent. For the proof we refer to [5].

Theorem 2.5. *Let $\mathfrak{M}$ be a closed O^* -algebra $\mathcal{D}$ with $\mathcal{D}$ a Fréchet domain of $\mathfrak{M}$. Assume that $\mathfrak{M}'_{\mathfrak{w}}\mathcal{D} \subset \mathcal{D}$. Let E' be a projection in $\mathfrak{M}'_{\mathfrak{w}}$ and assume that $\langle \mathfrak{M}'_{\mathfrak{w}}E'\mathcal{D} \rangle$ is essentially self-adjoint for $\mathfrak{M}''_{\mathfrak{w}c}$. Then, $(\mathfrak{M}_{E'})''_{\mathfrak{w}c} = (\mathfrak{M}''_{\mathfrak{w}c})_{E'}$.*

Proof. Since $\mathcal{D}[t_{\mathfrak{M}}]$ is a Fréchet space, the topology $t_{\mathfrak{M}}$ is defined by a sequence $\{T_n\}$ of elements of $\mathfrak{M}$ satisfying

$$\|\xi\| \leq \|T_1\xi\| \leq \|T_2\xi\|, \dots, \quad \forall \xi \in \mathcal{D}.$$

Any $X \in \mathcal{L}^\dagger(\mathcal{D})$ is a closed linear operator on the Fréchet space $\mathcal{D}[t_{\mathfrak{M}}]$ into the Hilbert space $\mathcal{H}$, which implies, by the closed graph theorem, that the topology $t_{\mathfrak{M}}$ is equivalent to the graph topology $t_{\mathcal{L}^\dagger(\mathcal{D})}$ defined by $\mathcal{L}^\dagger(\mathcal{D})$. This in turn implies that, for every $X \in \mathcal{L}^\dagger(\mathcal{D})$ there exist $n_0 \in \mathbb{N}$ and $\gamma > 0$ such that

$$(2) \quad \|X\xi\| \leq \gamma\|T_{n_0}\xi\|, \quad \forall \xi \in \mathcal{D}.$$

Take now an arbitrary $X \in (\mathfrak{M}_{E'})''_{\text{wc}}$. Since $XE' \in \mathcal{L}^\dagger(\mathcal{D})$, it follows from (2) that

$$(3) \quad \|XE'\xi\| \leq \gamma\|T_{n_0}\xi\|, \quad \forall \xi \in \mathcal{D}.$$

Since $\mathfrak{M}'_{\text{w}}$ is a von Neumann algebra, for every $C \in \mathfrak{M}'_{\text{w}}$ and $\xi \in \mathcal{D}$, we have

$$\begin{aligned} \|X_e(CE'\xi)\|^2 &= \|CXE'\xi\|^2 \\ &= \langle E'C^*CE'XE'\xi | XE'\xi \rangle \\ &= \|(E'C^*CE')^{1/2}XE'\xi\|^2 \\ &= \|XE'(E'C^*CE')^{1/2}\xi\|^2 \\ &\leq \gamma^2\|T_{n_0}(E'C^*CE')^{1/2}\xi\|^2 \\ &= \gamma^2\|(E'C^*CE')^{1/2}T_{n_0}\xi\|^2 \\ &= \gamma^2\|CE'T_{n_0}\xi\|^2 \\ &= \gamma^2\|T_{n_0}CE'\xi\|^2. \end{aligned}$$

Hence, the condition (i) of Lemma 2.3 is satisfied. By the same Lemma we then get the equality $(\mathfrak{M}_{E'})''_{\text{wc}} = (\mathfrak{M}'_{\text{wc}})_{E'}$. $\square$

Lemma 2.6. *Let $\mathfrak{M}$ be a closed O^* -algebra on a Fréchet domain $\mathcal{D}$ in $\mathcal{H}$ such that $t_{\mathfrak{M}}$ is defined by a sequence $\{T_n\}$ of essentially self-adjoint operators of $\mathfrak{M}$ whose spectral projections belong to $\mathfrak{M}$.*

Let $\mathcal{M}$ be an $\mathfrak{M}$ -invariant subspace of $\mathcal{D}$ such that the projection $P_{\mathcal{M}}$ onto $\overline{\mathcal{M}}$ belongs to $\mathfrak{M}'_{\text{w}}$.

Then $\mathcal{M}$ is essentially self-adjoint for $\mathfrak{M}$ or, equivalently, $P_{\mathcal{M}}\mathcal{D} = \overline{\mathcal{M}}^{t_{\mathfrak{M}}}$.

Proof. First we observe that $\overline{\mathcal{M}}^{t_{\mathfrak{M}}} \subset P_{\mathcal{M}}\mathcal{D}$. Hence we only need to show that the converse inclusion also holds. Take an arbitrary $\xi \in \mathcal{D}$. Then there exists a sequence $\{\xi_n\}$ in $\mathcal{M}$ such that $\lim_{n \rightarrow \infty} \|\xi_n - P_{\mathcal{M}}\xi\| = 0$. By (3), for every $X \in \mathfrak{M}$, there exists $n_0 \in \mathbb{N}$ and $\gamma > 0$ such that

$$(4) \quad \|X\xi\| \leq \gamma\|T_{n_0}\xi\|, \quad \forall \xi \in \mathcal{D}.$$

By the assumption, $\overline{T_{n_0}}$ is self-adjoint. Let $\overline{T_{n_0}} = \int_{-\infty}^{\infty} \lambda dE_{T_{n_0}}(\lambda)$ be the spectral resolution of $\overline{T_{n_0}}$. We put $E_k := E_{T_{n_0}}(k) - E_{T_{n_0}}(-k)$, $k \in \mathbb{N}$. Then, by assumption $E_k \upharpoonright \mathcal{D} \in \mathfrak{M}$, $\forall k \in \mathbb{N}$ and so, $E_k\xi_n \in \mathcal{M}$, $\forall k, n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} \|E_k\xi_n - E_k P_{\mathcal{M}}\xi\| = 0$. Furthermore, by (4) we have

$$\|XE_k\xi_n - XE_k\xi_m\| \leq \gamma\|T_{n_0}E_k(\xi_n - \xi_m)\| \rightarrow 0, \text{ as } n, m \rightarrow \infty.$$

Hence $E_k P_{\mathcal{M}}\xi \in \overline{D(X \upharpoonright \mathcal{M})}$.

Furthermore,

$$\begin{aligned} \|XE_k P_{\mathcal{M}}\xi - X P_{\mathcal{M}}\xi\| &\leq \gamma \|T_{n_0} E_k P_{\mathcal{M}}\xi - T_{n_0} P_{\mathcal{M}}\xi\| \\ &= \gamma \|P_{\mathcal{M}}(E_k T_{n_0}\xi - T_{n_0}\xi)\| \\ &\leq \gamma \|E_k T_{n_0}\xi - T_{n_0}\xi\| \rightarrow 0 \text{ as } k \rightarrow \infty, \end{aligned}$$

which implies that $P_{\mathcal{M}}\xi \in \overline{D(X \upharpoonright \mathcal{M})}$. Hence,

$$P_{\mathcal{M}}\xi \in \bigcap_{X \in \mathfrak{M}} \overline{D(X \upharpoonright \mathcal{M})} = \overline{\mathcal{M}}^{t_{\mathfrak{M}}}.$$

Thus $P_{\mathcal{M}}\mathcal{D} = \overline{\mathcal{M}}^{t_{\mathfrak{M}}}$. This is equivalent to $\mathcal{M}$ being essentially self-adjoint for $\mathfrak{M}$, since $\mathfrak{M}$ is clearly self-adjoint. $\square$

By Theorem 2.5 and Lemma 2.6 we have the following

Theorem 2.7. *Let $\mathfrak{M}$ be a closed O^* -algebra such that $t_{\mathfrak{M}}$ is defined by a sequence $\{T_n\}$ of essentially self-adjoint operators of $\mathfrak{M}$ whose spectral projections leave the domain $\mathcal{D}$ invariant. Let E' be a projection in $\mathfrak{M}'_w$. Then $(\mathfrak{M}_{E'})''_{wc} = (\mathfrak{M}''_{wc})_{E'}$.*

Proof. It can be shown that $\mathfrak{M}$ is a self-adjoint O^* -algebra on the Fréchet domain $\mathcal{D}$, so that $\mathfrak{M}'_w \mathcal{D} \subset \mathcal{D}$, $t_{\mathfrak{M}'_{wc}} = t_{\mathfrak{M}}$ and $\mathfrak{M}''_{wc}$ is a self-adjoint O^* -algebra on the Fréchet domain $\mathcal{D}$. Furthermore

$$P_{\langle \mathcal{M}'_w E' \mathcal{D} \rangle} = Z \in \mathfrak{M}'_w \cap (\mathfrak{M}'_w)' = (\mathfrak{M}''_{wc})'_w \cap ((\mathfrak{M}''_{wc})'_w)'.$$

Moreover, since the spectral projections of the operators T_n leave the domain $\mathcal{D}$ invariant, they automatically belong to $\mathfrak{M}''_{wc}$. Hence, the self-adjoint O^* -algebra $\mathfrak{M}''_{wc}$ and the $\mathfrak{M}''_{wc}$ -invariant subspace $\langle \mathcal{M}'_w E' \mathcal{D} \rangle$ of $\mathcal{D}$ satisfy all the conditions in Lemma 2.6, and so $Z\mathcal{D} = \overline{\langle \mathcal{M}'_w E' \mathcal{D} \rangle}^{t_{\mathfrak{M}''_{wc}}}$. Then, by Theorem 2.5, $(\mathfrak{M}_{E'})''_{wc} = (\mathfrak{M}''_{wc})_{E'}$. $\square$

Corollary 2.8. *Let T be an essentially self-adjoint operator in $\mathcal{H}$ and $\mathfrak{M}$ be a self-adjoint O^* -algebra on $\mathcal{D}^\infty(\overline{T}) := \bigcap_{n \in \mathbb{N}} D(\overline{T}^n)$, containing T . Let E' be a projection in $\mathfrak{M}'_w$. Then $(\mathfrak{M}_{E'})''_{wc} = (\mathfrak{M}''_{wc})_{E'}$.*

Proof. The spectral projections $E_T(\lambda)$, $\lambda \in \mathbb{R}$, of $\overline{T}$ satisfy:

- $E_T(\lambda) \in (\mathfrak{M}'_w)'$, $\forall \lambda \in \mathbb{R}$;
- $E_T(\lambda)\mathcal{H} \subset \mathcal{D}^\infty(\overline{T})$, $\forall \lambda \in \mathbb{R}$.

The statement then follows from Theorem 2.7. $\square$

Apart from GW^* -algebras, another unbounded generalization of von Neumann algebras is provided by the notion of *extended W^* -algebras*, shortly EW^* -algebras, defined as follows: A closed O^* -algebra $\mathfrak{M}$ on $\mathcal{D}$ is said to be an *EW*-algebra* if $(I + X^\dagger X)^{-1}$ exists in $\mathfrak{M}_b := \{A \in \mathfrak{M} : \overline{A} \in \mathcal{B}(\mathcal{H})\}$, for every $X \in \mathfrak{M}$ and $\overline{\mathfrak{M}}_b := \{\overline{A} \in \mathfrak{M} : A \in \mathfrak{M}_b\}$ is a von Neumann algebra [2]. It is easily shown that every EW^* -algebra on a Fréchet domain satisfies the conditions of Theorem 2.7 (remind that every symmetric element of an EW^* -algebra is essentially self-adjoint). Hence we have the following

Corollary 2.9. *Let $\mathfrak{M}$ be an EW^* -algebra on the Fréchet domain $\mathcal{D}$ in Hilbert space $\mathcal{H}$ and E' a projection in $\mathfrak{M}'_w$. Then, $(\mathfrak{M}_{E'})''_{wc} = (\mathfrak{M}''_{wc})_{E'}$.*

3. APPLICATIONS

In this section we show how to use the results of Section 2 in the analysis of the existence of conditional expectations for O^* -algebras, which were first studied in [3, 4].

Let $\mathfrak{M}$ be a given O^* -algebra on $\mathcal{D}$ in $\mathcal{H}$ with a strongly cyclic vector ξ_0 . Here $\xi_0 \in \mathcal{D}$ is said to be strongly cyclic for $\mathfrak{M}$ if $\overline{\mathfrak{M}\xi_0}^{\mathfrak{M}} = \mathcal{D}$. With no loss of generality we will assume that $\|\xi_0\| = 1$. Let $\mathfrak{N}$ be a O^* -subalgebra of $\mathfrak{M}$. A map $\mathcal{E}$ of $\mathfrak{M}$ onto $\mathfrak{N}$ is said to be a conditional expectation of $(\mathfrak{M}, \xi_0)$ w.r.t. $\mathfrak{N}$ if it satisfies the following conditions:

- (i) $\mathcal{E}(X)^\dagger = \mathcal{E}(X^\dagger)$, $\forall X \in \mathfrak{M}$, and $\mathcal{E}(A) = A$, $\forall A \in \mathfrak{N}$;
- (ii) $\mathcal{E}(XA) = \mathcal{E}(X)A$ and $\mathcal{E}(AX) = A\mathcal{E}(X)$, $\forall X \in \mathfrak{M}$, $\forall A \in \mathfrak{N}$;
- (iii) $\omega_{\xi_0}(\mathcal{E}(X)) = \omega_{\xi_0}(X)$, $\forall X \in \mathfrak{M}$, where ω_{ξ_0} is a state on $\mathfrak{M}$ defined as $\omega_{\xi_0}(X) = \langle X\xi_0 | \xi_0 \rangle$, $X \in \mathfrak{M}$.

In the case of von Neumann algebras Takesaki [7] characterized the existence of conditional expectations using Tomita's modular theory. Thus a conditional expectation does not necessarily exist for a general von Neumann algebra. Hence, in [3, 4] Ogi, Takakura and one of us considered a linear map $\mathcal{E}$ of a $\dagger$ -invariant subspace $D(\mathcal{E})$ of $\mathfrak{M}$ onto $\mathfrak{N}$ satisfying the above conditions (i), (ii) and (iii) on $D(\mathcal{E})$. Such a map is called an *unbounded conditional expectation* of $(\mathfrak{M}, \xi_0)$ w.r.t. $\mathfrak{N}$, and it was shown that there exists the largest unbounded conditional expectation $\mathcal{E}_{\mathfrak{N}}$ of $(\mathfrak{M}, \xi_0)$ w.r.t. $\mathfrak{N}$. Furthermore, the existence of conditional expectation was characterized, using Takesaki's result in the case of von Neumann algebras. But a deeper analysis is needed since, at that stage the problem was not solved even in the case of GW^* -algebras. One of the reason is that the reduction of a GW^* -algebra is not necessarily a GW^* -algebra, in contrast with the case of von Neumann algebras. For $\mathcal{E}_{\mathfrak{N}}$ to be a conditional expectations of $(\mathfrak{M}, \xi_0)$ w.r.t. $\mathfrak{N}$ (i.e., everywhere defined on $\mathfrak{M}$), the following result given in [4, Corollary 6.2] holds.

Lemma 3.1. *Let $\mathfrak{M}$ be a closed O^* -algebra on $\mathcal{D}$ in $\mathcal{H}$ such that $\mathfrak{M}'_w \mathcal{D} \subset \mathcal{D}$. Let ξ_0 be a strongly cyclic and separating vector, in the sense that $\overline{\mathfrak{M}'_w \xi_0} = \mathcal{H}$. Suppose that $\mathfrak{N}$ is a closed O^* -subalgebra of $\mathfrak{M}$ satisfying*

- (i) $\mathfrak{N}'_w \mathcal{D} \subset \mathcal{D}$;
 - (ii) $\mathfrak{N}\xi_0$ is essentially self-adjoint for $\mathfrak{N}$;
 - (iii) $\Delta''_{\xi_0}{}^{it}(\mathfrak{N}'_w)' \Delta''_{\xi_0}{}^{-it} = (\mathfrak{N}'_w)', \forall t \in \mathbb{R}$, where Δ''_{ξ_0} is the modular operator of the left Hilbert algebra $(\mathfrak{M}'_w)' \xi_0$;
 - (iv) $\mathfrak{N}_{P_{\mathfrak{N}}}$ is a GW^* -algebra on $P_{\mathfrak{N}}\mathcal{D}$, where $P_{\mathfrak{N}}$ is a projection of $\mathcal{H}$ onto $\overline{\mathfrak{N}\xi_0}$.
- Then $\mathcal{E}_{\mathfrak{N}}$ is a conditional expectation of $(\mathfrak{M}, \xi_0)$ w.r.t. $\mathfrak{N}$.*

By this Lemma and Theorem 2.5 we deduce the following

Theorem 3.2. *Let $\mathfrak{M}$ be a closed O^* -algebra on $\mathcal{D}$ in $\mathcal{H}$ with a strongly cyclic and separating vector ξ_0 such that $\mathfrak{M}'_w \mathcal{D} \subset \mathcal{D}$ and let $\mathfrak{N}$ be an O^* -subalgebra of $\mathfrak{M}$.*

Suppose that $\mathfrak{N}$ is a GW^ -algebra on a Fréchet domain $\mathcal{D}$ satisfying*

- (i) $\mathfrak{N}\xi_0$ and $\langle \mathfrak{N}'_w P_{\mathfrak{N}} \mathcal{D} \rangle$ are essentially self-adjoint for $\mathfrak{N}$;
- (ii) $\Delta''_{\xi_0}{}^{it}(\mathfrak{N}'_w)' \Delta''_{\xi_0}{}^{-it} = (\mathfrak{N}'_w)', \forall t \in \mathbb{R}$.

Then $\mathcal{E}_{\mathfrak{N}}$ is a conditional expectation of $(\mathfrak{M}, \xi_0)$ w.r.t. $\mathfrak{N}$.

By Lemma 3.1 and Theorem 2.7 we deduce the following

Theorem 3.3. *Let $(\mathfrak{M}, \xi_0)$ be as in Theorem 3.2 and $\mathfrak{N}$ an O^* -subalgebra of $\mathfrak{M}$. Suppose $\mathfrak{N}$ is a GW^* -algebra on $\mathcal{D}$ satisfying:*

- (i) *$t_{\mathfrak{N}}$ is defined by a sequence $\{T_n\}$ of essentially self-adjoint operators in $\mathfrak{N}$ whose spectral projections leave the domain $\mathcal{D}$ invariant;*
- (ii) *$\Delta_{\xi_0}''^{it}(\mathfrak{N}'_w)' \Delta_{\xi_0}''^{-it} = (\mathfrak{N}'_w)', \forall t \in \mathbb{R}$.*

Then $\mathcal{E}_{\mathfrak{N}}$ is a conditional expectation of $(\mathfrak{M}, \xi_0)$ w.r.t. $\mathfrak{N}$.

Proof. By Lemma 2.6, $\mathfrak{N}\xi_0$ and $\langle \mathfrak{N}'_w P_{\mathfrak{N}} \mathcal{D} \rangle$ are essentially self-adjoint for $\mathfrak{N}$; by Theorem 3.2 it follows that $\mathcal{E}_{\mathfrak{N}}$ is a conditional expectation of $(\mathfrak{M}, \xi_0)$ w.r.t. $\mathfrak{N}$. $\square$

Corollary 3.4. *Let $(\mathfrak{M}, \xi_0)$ be as in Theorem 3.2 and $\mathfrak{N}$ an O^* -subalgebra of $\mathfrak{M}$. Suppose $\mathfrak{N}$ is a GW^* -algebra on $\mathcal{D}^\infty(\overline{T})$ where T is an essentially self-adjoint operator in $\mathfrak{N}$, and $\Delta_{\xi_0}''^{it}(\mathfrak{N}'_w)' \Delta_{\xi_0}''^{-it} = (\mathfrak{N}'_w)', \forall t \in \mathbb{R}$. Then, $\mathcal{E}_{\mathfrak{N}}$ is a conditional expectation of $(\mathfrak{M}, \xi_0)$ w.r.t. $\mathfrak{N}$.*

Corollary 3.5. *Let $(\mathfrak{M}, \xi_0)$ be as in Theorem 3.2 and $\mathfrak{N}$ an O^* -subalgebra of $\mathfrak{M}$. Suppose $\mathfrak{N}$ is an EW^* -algebra on a Fréchet domain $\mathcal{D}$ satisfying $\Delta_{\xi_0}''^{it}(\mathfrak{N}'_w)' \Delta_{\xi_0}''^{-it} = (\mathfrak{N}'_w)', \forall t \in \mathbb{R}$. Then $\mathcal{E}_{\mathfrak{N}}$ can be extended to a conditional expectation $\mathcal{E}_{\mathfrak{N}'_{wc}}$ of $(\mathfrak{M}'_{wc}, \xi_0)$ w.r.t. $\mathfrak{N}'_{wc}$.*

Acknowledgement One of us (A.I.) acknowledges the hospitality of the Dipartimento di Matematica ed Applicazioni, Università di Palermo, where this work was performed.

REFERENCES

- [1] J. Dixmier, *Von Neumann algebras*, North-Holland Publ. Comp., Amsterdam, 1981.
- [2] A. Inoue, *On a class of unbounded operator algebras*, Pacific J. Math. **65**, 11-95 (1976).
- [3] A. Inoue, H. Ogi, M. Takakura, *Conditional expectations for unbounded operator algebras*, Int. J. Math. Math. Sci. 2007, Art. ID 80152, 22 pp.
- [4] A. Inoue, H. Ogi, M. Takakura, *Conditional expectations for O^* -algebras*, Contemporary Math., **427**, 225-234 (2007)
- [5] R.T. Powers, *Self-adjoint algebras of unbounded operators*, Commun. Math. Phys., **21**, 85-124 (1971).
- [6] K. Schmüdgen, *Unbounded Operator Algebras and Representation Theory*, Birkhäuser-Verlag, Basel, 1990.
- [7] M. Takesaki, *Conditional expectations in a von Neumann algebra*, J. Funct. Anal., **9**, 306-327 (1972).

DIPARTIMENTO DI METODI E MODELLI MATEMATICI, FACOLTÀ D'INGEGNERIA, UNIVERSITÀ DI PALERMO, I-90128 PALERMO, ITALY

E-mail address: bagarell@unipa.it

DEPARTMENT OF APPLIED MATHEMATICS, FUKUOKA UNIVERSITY, 814-0180, JAPAN

E-mail address: a-inoue@fukuoka-u.ac.jp

DIPARTIMENTO DI MATEMATICA ED APPLICAZIONI, UNIVERSITÀ DI PALERMO, I-90123 PALERMO, ITALY

E-mail address: trapani@unipa.it